

IDENTITIES FOR HYPERGEOMETRIC INTEGRALS OF DIFFERENT DIMENSIONS

V. TARASOV^{*} AND A. VARCHENKO[◊]

*St. Petersburg Branch of Steklov Mathematical Institute
Fontanka 27, St. Petersburg 191011, Russia*

[◊] *Department of Mathematics, University of North Carolina at Chapel Hill
Chapel Hill, NC 27599, USA*

May, 2003

ABSTRACT. Given complex numbers m_1, l_1 and nonnegative integers m_2, l_2 , such that $m_1 + m_2 = l_1 + l_2$, we define $\hat{l}_2$ -dimensional hypergeometric integrals $I_{a,b}(z; m_1, m_2, l_1, l_2)$, $a, b = 0, \dots, \min(m_2, l_2)$, depending on a complex parameter z . We show that $I_{a,b}(z; m_1, m_2, l_1, l_2) = I_{a,b}(z; l_1, l_2, m_1, m_2)$, thus establishing an equality of $\hat{l}_2$ and $\hat{m}_2$ -dimensional integrals. This identity allows us to study asymptotics of the integrals with respect to their dimension in some examples. The identity is based on the $(\mathfrak{gl}_k, \mathfrak{gl}_n)$ duality for the KZ and dynamical differential equations.

1. INTRODUCTION

Let κ be a positive number. Let m_1, l_1 be complex numbers and m_2, l_2 nonnegative integers such that

$$m_1 + m_2 = l_1 + l_2.$$

We say that an integer a is *admissible* with respect to m_2, l_2 if

$$0 \leq a \leq \min(m_2, l_2).$$

For a pair of admissible numbers a, b we define a function $I_{a,b}(z; m_1, m_2, l_1, l_2)$ of one complex variable z . The function is defined as an $\hat{l}_2$ -dimensional hypergeometric integral of the form

$$(1.1) \quad \begin{aligned} I_{a,b}(z; m_1, m_2, l_1, l_2) &= \\ &= C_b(m_1, m_2, l_1, l_2) \int_{\gamma_{l_2, b}(z)} \Phi_{l_2}(t, z; m_1, m_2)^{1/\kappa} w_{l_2-a, a}(t, z) dt^{l_2}. \end{aligned}$$

Here $t = (t_1, \dots, t_{l_2})$, $dt^{l_2} = dt_1 \wedge \dots \wedge dt_{l_2}$. The constant $C_b(m_1, m_2, l_1, l_2)$ and the functions $\Phi_{l_2}(t, z; m_1, m_2)$ and $w_{l_2-a, a}(t, z)$ are defined below. The $\hat{l}_2$ -dimensional integration contour $\gamma_{l_2, b}(z)$ lies in $\mathbb{C}^{l_2}$ and is also defined below.

^{*}Supported in part by RFFI grant 02-01-00085a and CRDF grant RM1-2334-02

E-mail: vt@pdmi.ras.ru

[◊]Supported in part by NSF grant DMS-0244579

E-mail: anv@email.unc.edu


The *master* function Φ_l is defined by the formula

$$(1.2) \quad \Phi_l(t_1, \dots, t_l, z; m_1, m_2) = e^{-\sum_{u=1}^l t_u} \prod_{u=1}^l (-t_u)^{-m_1} (z - t_u)^{-m_2} \prod_{1 \leq u < v \leq l} (t_u - t_v)^2.$$

The *weight* function $w_{l-a,a}$ is defined by the formula

$$(1.3) \quad w_{l-a,a}(t_1, \dots, t_l, z) = \text{Sym} \left[\prod_{u=1}^{l-a} \frac{1}{-t_u} \prod_{u=l-a+1}^l \frac{1}{z - t_u} \right]$$

where $\text{Sym } f(t_1, \dots, t_l) = \sum_{\sigma \in S_l} f(t_{\sigma_1}, \dots, t_{\sigma_l})$. We define the integral in (1.1) by analytic continuation from the region $\text{Im } z > 0$. For $\text{Im } z > 0$ the integration contour $\gamma_{l,b}(z)$ is shown in the picture.



Picture 1. The contour $\gamma_{l,b}(z)$.

It has the form

$$\gamma_{l,b}(z) = \{ (t_1, \dots, t_l) \in \mathbb{C}^l \mid t_u \in \mathcal{C}_u, \ u = 1, \dots, l \}.$$

Here $\mathcal{C}_u$, $u = 1, \dots, l$, are non-intersecting oriented loops in $\mathbb{C}$. The first b loops start at $+\infty$, go around z , and return to $+\infty$. For $1 \leq u < v \leq b$, the loop $\mathcal{C}_u$ lies inside the loop $\mathcal{C}_v$. The last $l - b$ loops start at $+\infty$, go around 0 , and return to $+\infty$. For $b + 1 \leq u < v \leq l$, the loop $\mathcal{C}_u$ lies inside the loop $\mathcal{C}_v$.

If $\text{Im } z > 0$, then we fix a univalued branch of $\Phi_l(t, z; m_1, m_2)^{1/\kappa}$ over $\gamma_{l,b}(z)$ by fixing the arguments of all factors of Φ_l . Namely, we assume that at the point of $\gamma_{l,b}(z)$ where all numbers $z - t_1, \dots, z - t_b, -t_{b+1}, \dots, -t_l$ are positive we have

$$\begin{aligned} \arg(-t_u) &\in (-\pi, 0) \text{ for } u = 1, \dots, b, \text{ and } \arg(-t_u) = 0 \text{ for } u = b + 1, \dots, l; \\ \arg(z - t_u) &= 0 \text{ for } u = 1, \dots, b, \text{ and } \arg(z - t_u) \in (0, \pi) \text{ for } u = b + 1, \dots, l; \\ \arg(t_u - t_v) &= 0 \text{ for } u < v \leq b \text{ and for } b < u < v; \\ \arg(t_u - t_v) &\in (0, \pi) \text{ for } u \leq b < v. \end{aligned}$$

Set

$$\begin{aligned} C_b(m_1, m_2, l_1, l_2) &= \kappa^{(l_1+1)l_2/\kappa} e^{-\pi i b l_2/\kappa} (2i \Gamma(-1/\kappa))^{-l_2} \times \\ &\times \prod_{j=0}^{b-1} \frac{1}{\sin(\pi(j+1)/\kappa)} \prod_{j=0}^{l_2-b-1} \frac{1}{\sin(\pi(j+1)/\kappa)} \prod_{j=0}^{l_2-1} \frac{\Gamma(1 + (m_1 - j)/\kappa)}{\Gamma(1 + (j+1)/\kappa)}. \end{aligned}$$

Theorem 1. Let m_1, l_1 be complex numbers and m_2, l_2 nonnegative integers, such that $m_1 + m_2 = l_1 + l_2$. Let κ be a generic positive number. Then for any a, b , admissible with respect to m_2, l_2 , we have

$$I_{a,b}(z; m_1, m_2, l_1, l_2) = I_{a,b}(z; l_1, l_2, m_1, m_2).$$

This theorem claims that an $\hat{l}_2$ -dimensional integral equals an $\hat{m}_2$ -dimensional integral. Note that in the first integral the numbers m_1, m_2 are exponents of its master function, while in the second integral m_2 is its dimension and m_1 is not present explicitly.

It is well known that studying asymptotics of integrals with respect to their dimension is an interesting problem appearing, for instance, in the theory of orthogonal polynomials and in the matrix models. The duality of the theorem allows us to study asymptotics of integrals with respect to their dimension in some examples. Namely, assume that in the 4-tuple (m_1, m_2, l_1, l_2) the nonnegative integer l_2 tends to infinity, while the numbers m_1 and m_2 remain fixed. Then $I_{a,b}(z; m_1, m_2, l_1, l_2)$ is an integral of growing dimension l_2 , whose master function has fixed exponents m_1, m_2 . At the same time, $I_{a,b}(z; l_1, l_2, m_1, m_2)$ is an integral of the fixed dimension m_2 , whose master function has growing exponents l_1, l_2 . The asymptotics of $I_{a,b}(z; l_1, l_2, m_1, m_2)$ can be calculated using the steepest descent method, see a more precise statement below.

To prove Theorem 1 we show that the matrices:

$$(I_{a,b}(z; m_1, m_2, l_1, l_2))_{0 \leq a, b \leq \min(m_2, l_2)} \quad \text{and} \quad (I_{a,b}(z; l_1, l_2, m_1, m_2))_{0 \leq a, b \leq \min(m_2, l_2)}$$

satisfy the same first order linear differential equation with respect to z , and have the same asymptotics as z tends to infinity, see Lemmas 2 and 3. Hence, the matrices are equal.

The fact that both matrices satisfy the same differential equation is based on the duality of the KZ and dynamical differential equations for $\mathfrak{gl}_k$ and $\mathfrak{gl}_n$ [TV1]. Namely, according to [FMTV] the KZ and dynamical differential equations for $\mathfrak{gl}_k$ have the hypergeometric solutions, and the KZ and dynamical differential equations for $\mathfrak{gl}_n$ also have the hypergeometric solutions. The $(\mathfrak{gl}_k, \mathfrak{gl}_n)$ duality allows one to conclude that both sets of solutions solve the same differential equations. Thus, the two sets of solutions, in principle, can be identified. Theorem 1 is a realization of this idea for $k = n = 2$. We will discuss the case of an arbitrary pair k, n in a separate paper.

In Section 2 we give a proof of Theorem 1. In Section 3 we discuss the $(\mathfrak{gl}_2, \mathfrak{gl}_2)$ duality for the KZ and dynamical differential equations. In Section 4 we construct hypergeometric solutions of the KZ and dynamical differential equations and relate them to the integrals $I_{a,b}(z; m_1, m_2, l_1, l_2)$. In Section 5 we calculate an example of asymptotics of hypergeometric integrals with respect to their dimension.

The authors thank Y. Markov for numerous useful discussions.

2. PROOF OF THEOREM 1

Introduce matrices A and $B(m_1, m_2, l_1, l_2)$ with entries

$$A_{a,b} = a \delta_{a,b},$$

$$\begin{aligned} B_{a,b}(m_1, m_2, l_1, l_2) = & (2a^2 - a(2l_2 + m_2 - m_1) + m_2 l_2) \delta_{a,b} + \\ & + a(l_2 - m_1 - a) \delta_{a-1,b} - (m_2 - a)(l_2 - a) \delta_{a+1,b}, \end{aligned}$$

$$a, b = 0, \dots, \min(m_2, l_2).$$

Lemma 2. *The matrix $\hat{I}(z; m_1, m_2, l_1, l_2) = (I_{a,b}(z; m_1, m_2, l_1, l_2))_{0 \leq a, b \leq \min(m_2, l_2)}$ obeys the ordinary differential equation*

$$(2.1) \quad \left(\kappa \frac{\partial}{\partial z} + \frac{B(m_1, m_2, l_1, l_2)}{z} + A \right) \hat{I}(z; m_1, m_2, l_1, l_2) = 0.$$

The lemma admits a straightforward proof using integration by parts. In Section 4 we give another proof of Lemma 2 using the hypergeometric solutions of the *KZ* and dynamical differential equations.


Lemma 3. *Fix a positive number ε less than $\pi/2$. If z tends to infinity inside the sector $\varepsilon < \arg z < \pi - \varepsilon$, then*

$$I_{a,b}(z; m_1, m_2, l_1, l_2) = e^{-bz/\kappa} (\kappa/z)^{(2b^2 - b(2l_2 + m_2 - m_1) + m_2 l_2)/\kappa} e^{\pi i b(m_1 - l_2)/\kappa} \times \\ \times \prod_{j=0}^{b-1} \frac{\Gamma(1 + (j+1)/\kappa) \Gamma(1 + (m_1 - l_2 + j + 1)/\kappa)}{\Gamma(1 + (m_2 - j)/\kappa) \Gamma(1 + (l_2 - j)/\kappa)} (\delta_{a,b} + O(z^{-1})).$$

Proof. For a positive number κ , complex number m , and a nonnegative integer l , consider the Selberg type integral

$$J_l(m) = \int_{\delta_l} e^{-\sum_{u=1}^l s_u/\kappa} \prod_{u=1}^l (-s_u)^{-1-m/\kappa} \prod_{1 \leq u < v \leq l} (s_u - s_v)^{2/\kappa} ds^k.$$

The integration contour δ_l has the form $\delta_l = \{(s_1, \dots, s_l) \in \mathbb{C}^l \mid s_u \in \mathcal{C}_u, u = 1, \dots, l\}$, see the picture.



Picture 2. The contour δ_l .

Here $\mathcal{C}_u$, $u = 1, \dots, l$, are non-intersecting oriented loops in $\mathbb{C}$. The loops start at $+\infty$, go around 0, and return to $+\infty$. For $u < v$, the loop $\mathcal{C}_u$ lies inside the loop $\mathcal{C}_v$. We fix a univalued branch of the integrand by assuming that at the point of δ_l where all numbers $-s_1, \dots, -s_l$ are positive we have $\arg(-s_u) = 0$ for $u = 1, \dots, l$, and $\arg(s_u - s_v) = 0$ for $1 \leq u < v \leq l$.

The formula for $J_l(m)$ is well known:

$$(2.2) \quad J_l(m) = \kappa^{l(l-1-m)/\kappa} \prod_{j=0}^{l-1} \frac{-2\pi i \Gamma(1 - 1/\kappa)}{\Gamma(1 + (m - j)/\kappa) \Gamma(1 - (j + 1)/\kappa)},$$

for example, cf. [TV2], [MTV].

It is easy to see that

$$I_{a,b}(z; m_1, m_2, l_1, l_2) = e^{-bz/\kappa} z^{-(2b^2 - b(2l_2 + m_2 - m_1) + m_2 l_2)/\kappa} e^{\pi i b m_1/\kappa} \times \\ \times (l_2 - b)! b! C_b(m_1, m_2, l_1, l_2) J_{l_2-b}(m_1) J_b(m_2) (\delta_{a,b} + O(z^{-1})).$$

Substituting formulas for $C_b(m_1, m_2, l_1, l_2)$ and $J_l(m)$ we get the lemma. $\square$

Proof of Theorem 1. By Lemma 2, since $B(m_1, m_2, l_1, l_2) = B(l_1, l_2, m_1, m_2)$, the matrices $\hat{I}(z; m_1, m_2, l_1, l_2)$ and $\hat{I}(z; l_1, l_2, m_1, m_2)$ obey the same first order ordinary differential equation. Therefore,

$$\hat{I}(z; m_1, m_2, l_1, l_2) = \hat{I}(z; l_1, l_2, m_1, m_2) X(l_1, l_2, m_1, m_2).$$

Lemma 3 implies that the connection matrix X is simultaneously lower and upper triangular with the unit diagonal. Hence X is the identity. Theorem 1 is proved. $\square$

3. THE $(\mathfrak{gl}_2, \mathfrak{gl}_2)$ DUALITY FOR KZ AND DYNAMICAL DIFFERENTIAL EQUATIONS

3.1. The KZ and dynamical differential equations for $\mathfrak{gl}_2$. Let $E_{i,j}$, $i, j = 1, 2$, be the standard generators of the Lie algebra $\mathfrak{gl}_2$ over $\mathbb{C}$.

We have the root decomposition $\mathfrak{gl}_2 = \mathfrak{n}_+ \oplus \mathfrak{h} \oplus \mathfrak{n}_-$ where

$$\mathfrak{n}_+ = \mathbb{C} \cdot E_{1,2}, \quad \mathfrak{h} = \mathbb{C} \cdot E_{1,1} \oplus \mathbb{C} \cdot E_{2,2}, \quad \mathfrak{n}_- = \mathbb{C} \cdot E_{2,1}.$$

We identify the Cartan subalgebra $\mathfrak{h}$ with $\mathbb{C}^2$ mapping $\lambda = \lambda_1 E_{1,1} + \lambda_2 E_{2,2}$ to (λ_1, λ_2) .

The Casimir element $\Omega \in \mathfrak{gl}_2^{\otimes 2}$ is defined by the formula

$$\Omega = E_{1,1} \otimes E_{1,1} + E_{2,2} \otimes E_{2,2} + E_{1,2} \otimes E_{2,1} + E_{2,1} \otimes E_{1,2}.$$

Fix a non-zero complex number κ . Let $V_1, \dots, V_n$ be $\mathfrak{gl}_2$ -modules. Set $V = V_1 \otimes \dots \otimes V_n$. The KZ operators $\nabla_a(z_1, \dots, z_n, \lambda_1, \lambda_2, \kappa)$, $a = 1, \dots, n$, acting on V -valued functions of complex variables $z_1, \dots, z_n, \lambda_1, \lambda_2$ are defined by the formula

$$\nabla_a(z_1, \dots, z_n, \lambda_1, \lambda_2, \kappa) = \kappa \frac{\partial}{\partial z_a} - \sum_{\substack{b=1 \\ b \neq a}}^n \frac{\Omega^{(a,b)}}{z_a - z_b} - \lambda_1 E_{1,1}^{(a)} - \lambda_2 E_{2,2}^{(a)}, \quad a = 1, \dots, n.$$

The operator $\Omega^{(a,b)} : V \rightarrow V$ acts as Ω on $V_a \otimes V_b$, and as the identity on all other tensor factors. Similarly, the operator $E_{i,i}^{(a)}$ acts as $E_{i,i}$ on V_a and as the identity on all other tensor factors. The KZ equations for a function $U(z_1, \dots, z_n, \lambda_1, \lambda_2)$ are

$$\nabla_a(z_1, \dots, z_n, \lambda_1, \lambda_2, \kappa) U(z_1, \dots, z_n, \lambda_1, \lambda_2) = 0, \quad a = 1, \dots, n.$$

The dynamical differential operators $D_i(z_1, \dots, z_n, \lambda_1, \lambda_2, \kappa)$, $i = 1, 2$, acting on V -valued functions of complex variables $z_1, \dots, z_n, \lambda_1, \lambda_2$ are defined by the formulae

$$D_1(z_1, \dots, z_n, \lambda_1, \lambda_2, \kappa) = \kappa \frac{\partial}{\partial \lambda_1} - \frac{1}{\lambda_1 - \lambda_2} (E_{2,1} E_{1,2} - E_{2,2}) - \sum_{a=1}^n z_a E_{1,1}^{(a)},$$

$$D_2(z_1, \dots, z_n, \lambda_1, \lambda_2, \kappa) = \kappa \frac{\partial}{\partial \lambda_2} - \frac{1}{\lambda_2 - \lambda_1} (E_{2,1} E_{1,2} - E_{2,2}) - \sum_{a=1}^n z_a E_{2,2}^{(a)}.$$

The dynamical differential equations for a function $U(z_1, \dots, z_n, \lambda_1, \lambda_2)$ are

$$D_i(z_1, \dots, z_n, \lambda_1, \lambda_2, \kappa) U(z_1, \dots, z_n, \lambda_1, \lambda_2) = 0, \quad i = 1, 2.$$

Theorem 4 ([FMTV]). *The KZ equations and dynamical differential equations form a compatible system of equations. Namely, we have*

$$[\nabla_a, \nabla_b] = 0, \quad [\nabla_a, D_i] = 0 \quad [D_1, D_2] = 0,$$

for $a, b = 1, \dots, n$, and $i = 1, 2$.

The KZ and dynamical differential operators preserve the weight decomposition of V . Thus the KZ and dynamical differential equations can be considered as equations on a function $U(z_1, \dots, z_n, \lambda_1, \lambda_2)$ taking values in a given weight subspace of V .

3.2. The duality. For a complex number m , denote M_m the $\hat{\mathfrak{gl}}_2$ -Verma module with highest weight $(m, 0)$ and highest weight vector v_m . The vectors $E_{2,1}^d v_m$, $d \in \mathbb{Z}_{\geq 0}$, form a basis in M_m .

For a non-negative integer m , denote L_m the irreducible $\mathfrak{gl}_2$ -module with highest weight $(m, 0)$ and highest weight vector v_m . The vectors $E_{2,1}^d v_m$, $d = 0, \dots, m$, form a basis in L_m .

Let m_1, l_1 be complex numbers and m_2, l_2 non-negative integers such that $m_1 + m_2 = l_1 + l_2$. Consider the weight subspace $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$ of the tensor product $M_{m_1} \otimes L_{m_2}$. The weight subspace has a basis

$$(3.1) \quad F^a(m_1, m_2, l_1, l_2) = \frac{1}{(l_2 - a)! a!} E_{2,1}^{l_2 - a} v_{m_1} \otimes E_{2,1}^a v_{m_2}, \quad a = 0, \dots, \min(m_2, l_2).$$

There is a linear isomorphism

$$\varphi : (M_{m_1} \otimes L_{m_2})[l_1, l_2] \rightarrow (M_{l_1} \otimes L_{l_2})[m_1, m_2], \quad F^a(m_1, m_2, l_1, l_2) \mapsto F^a(l_1, l_2, m_1, m_2).$$

Theorem 5 ([TV1]). *Under the isomorphism φ the system of KZ and dynamical differential equations with values in $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$ turns into the system of dynamical differential and KZ equations with values in $(M_{l_1} \otimes L_{l_2})[m_1, m_2]$:*

$$\nabla_a(z_1, z_2, \lambda_1, \lambda_2, \kappa) \varphi = \varphi D_a(\lambda_1, \lambda_2, z_1, z_2, \kappa),$$

$$D_a(z_1, z_2, \lambda_1, \lambda_2, \kappa) \varphi = \varphi \nabla_a(\lambda_1, \lambda_2, z_1, z_2, \kappa),$$

for $a = 1, 2$.

4. HYPERGEOMETRIC SOLUTIONS

4.1. Construction of hypergeometric solutions. Define the *master* function by the formula

$$\begin{aligned} \Phi(s_1, \dots, s_{l_2}, z_1, z_2, \lambda_1, \lambda_2; m_1, m_2) &= \\ &= e^{\lambda_1(m_1 z_1 + m_2 z_2) - (\lambda_1 - \lambda_2) \sum_{u=1}^{l_2} s_u} (\lambda_1 - \lambda_2)^{-l_2} (z_1 - z_2)^{m_1 m_2} \times \\ &\times \prod_{u=1}^{l_2} (z_1 - s_u)^{-m_1} (z_2 - s_u)^{-m_2} \prod_{1 \leq u < v \leq l_2} (s_u - s_v)^2. \end{aligned}$$

For $0 \leq a \leq l_2$, define the *weight* function by the formula

$$w_{l_2-a,a}(s_1, \dots, s_{l_2}, z_1, z_2) = \text{Sym} \left[\prod_{u=1}^{l_2-a} \frac{1}{z_1 - s_u} \prod_{u=l_2-a+1}^{l_2} \frac{1}{z_2 - s_u} \right].$$

Define an $\hat{\mathfrak{gl}}_2$ $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$ -valued differential $\hat{\mathfrak{gl}}_2$ -form by the formula

$$\omega(s_1, \dots, s_{l_2}, z_1, z_2) = \sum_{a=0}^{l_2} w_{l_2-a,a}(s_1, \dots, s_{l_2}, z_1, z_2) ds^{l_2} \frac{1}{(l_2 - a)! a!} E_{2,1}^{l_2 - a} v_{m_1} \otimes E_{2,1}^a v_{m_2}.$$

Fix a complex number κ . Define an $\wedge(M_{m_1} \otimes L_{m_2})[l_1, l_2]$ -valued function by the formula

$$(4.1) \quad U_\delta(z_1, z_2, \lambda_1, \lambda_2) = \int_{\delta(z_1, z_2, \lambda_1, \lambda_2)} (\Phi(s_1, \dots, s_{l_2}, z_1, z_2, \lambda_1, \lambda_2; m_1, m_2))^{1/\kappa} \omega(s_1, \dots, s_{l_2}, z_1, z_2).$$

The function depends on the choice of integration chains $\delta(z_1, z_2, \lambda_1, \lambda_2)$. We assume that for each $z_1, z_2, \lambda_1, \lambda_2$ the chain lies in $\mathbb{C}^{l_2}$ with coordinates $s_1, \dots, s_{l_2}$, and the chains form a horizontal family of $\wedge l_2$ -dimensional homology classes with respect to the multivalued function $\Phi(s_1, \dots, s_{l_2}, z_1, z_2, \lambda_1, \lambda_2; m_1, m_2)^{1/\kappa}$, see a more precise statement below and in [FMTV].

Theorem 6. *For any choice of the horizontal family δ , the function $U_\delta(z_1, z_2, \lambda_1, \lambda_2)$ is a solution of the KZ and dynamical differential equations with values in $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$ and parameter κ .*

The theorem is a corollary of Theorem 3.1 in [FMTV].

The solutions of the KZ and dynamical differential equations constructed in this section are called the *hypergeometric solutions*.

4.2. Proof of Lemma 2. Consider the ordinary differential equation for a function $\Psi(x)$ taking values in $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$:

$$(4.2) \quad \left(\kappa \frac{\partial}{\partial x} - \frac{\Omega - m_1 m_2}{x} + E_{2,2}^{(2)} \right) \Psi(x) = 0.$$

One can directly check the following statement.

Lemma 7. *Let $\Psi(x)$ be a solution of equation (4.2). Then the function*

$$(4.3) \quad U(z_1, z_2, \lambda_1, \lambda_2) = e^{(z_1 \lambda_1 (m_1 - l_2) + z_1 \lambda_2 l_2 + z_2 \lambda_1 m_2)/\kappa} \times \\ \times (z_1 - z_2)^{m_1 m_2/\kappa} (\lambda_1 - \lambda_2)^{l_1 l_2/\kappa} \Psi(-(\lambda_1 - \lambda_2)(z_1 - z_2))$$

is a solution of the KZ and dynamical differential equations with values in $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$ and parameter κ . Moreover, formula (4.3) defines the one-to-one correspondence of solutions of the respective equations.

For any $b = 0, \dots, \min(m_2, l_2)$ set

$$\bar{I}_b(z; m_1, m_2, l_1, l_2) = \sum_{a=0}^{l_2} I_{a,b}(z; m_1, m_2, l_1, l_2) \frac{1}{(l_2 - a)! a!} E_{2,1}^{l_2 - a} v_{m_1} \otimes E_{2,1}^a v_{m_2}.$$

Here the actual range of summation is until $a = \min(m_2, l_2)$, since $E_{2,1}^a v_{m_2} = 0$ for $a > m_2$.

Lemma 8. *The function*

$$(4.4) \quad U_b(z_1, z_2, \lambda_1, \lambda_2) = e^{(z_1 \lambda_1 (m_1 - l_2) + z_1 \lambda_2 l_2 + z_2 \lambda_1 m_2)/\kappa} \times \\ \times (z_1 - z_2)^{m_1 m_2/\kappa} (\lambda_1 - \lambda_2)^{l_1 l_2/\kappa} \bar{I}_b(-(\lambda_1 - \lambda_2)(z_1 - z_2); m_1, m_2, l_1, l_2))$$

is a solution of the KZ and dynamical differential equations with values in $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$ and parameter κ .

Proof. Change the variables $t_1, \dots, t_{l_2}$ in the integrals $I_{a,b}(z; m_1, m_2, l_1, l_2)$ to the new variables

$$s_u = t_u(\lambda_1 - \lambda_2)^{-1} + z_1, \quad u = 1, \dots, l_2.$$

Then the right hand side of (4.4) takes the form of integral (4.1) with the chain $\delta(z_1, z_2, \lambda_1, \lambda_2)$ induced by the chain $\gamma_b(-(\lambda_1 - \lambda_2)(z_1 - z_2))$ under the described change of integration variables. One can see that δ is a horizontal family of chains. Therefore, the statement follows from Theorem 6. $\square$

To complete the proof of Lemma 2 observe that equation (4.2) written in basis (3.1) of the weight subspace $(M_{m_1} \otimes L_{m_2})[l_1, l_2]$ becomes equation (2.1). Hence, Lemma 7 yields Lemma 2.

5. REMARKS ON ASYMPTOTICS WITH RESPECT TO DIMENSION

Let κ be a positive number. Consider a 4-tuple m_1, m_2, l_1, l_2 of complex numbers such that m_2, l_2 are natural numbers and $m_1 + m_2 = l_1 + l_2$. For any integers a, b , admissible with respect to the pair m_2, l_2 , introduce an l_2 -dimensional integral $K_{a,b}(z; m_1, m_2, l_1, l_2)$ by the formula

$$K_{a,b}(z; m_1, m_2, l_1, l_2) = \int_{\gamma_{l_2,b}(z)} \Phi_{l_2}(t, z; m_1, m_2)^{1/\kappa} w_{l_2-a,a}(t, z) dt^{l_2},$$

where the cycle $\gamma_{l_2,b}(z)$ and the functions $\Phi_{l_2}(t, z; m_1, m_2)$, $w_{l_2-a,a}(t, z)$ are defined in the Introduction. Recall that $\gamma_{l_2,b}(z)$ is the $\hat{l}_2$ -dimensional multi-loop with b loops going around z and the remaining $l_2 - b$ loops going around 0.

Corollary 9. *We have*

$$K_{a,b}(z, m_1, m_2, l_1, l_2) = e^{\pi i b(l_2 - m_2)} \frac{(l_2 - b)! J_{l_2-b}(m_1) J_b(m_2)}{(m_2 - b)! J_{m_2-b}(l_1) J_b(l_2)} K_{a,b}(z, l_1, l_2, m_1, m_2).$$

Assume that the numbers m_1, m_2 are fixed and l_2 tends to infinity. Then the integral $K_{a,b}(z, m_1, m_2, l_1, l_2)$ is a hypergeometric integral of growing dimension l_2 with fixed exponents m_1, m_2 . In the right hand side we have $J_{l_2-b}(m_1)$, the Selberg integral of growing dimension; $J_{m_2-b}(l_1), J_b(l_2)$, the Selberg integrals of fixed dimension with growing exponents l_1, l_2 ; $K_{a,b}(z, l_1, l_2, m_1, m_2)$, the hypergeometric integral of fixed dimension m_2 with growing exponents l_1, l_2 .


The Selberg integrals are given explicitly by formula (2.2). The asymptotics of the integral $K_{a,b}(z, l_1, l_2, m_1, m_2)$ can be calculated by the steepest descent method.

Here is an example. If $m_2 = 1$, then the integral $K_{a,b}(z, l_1, l_2, m_1, m_2)$ is one-dimensional and its asymptotics are described as follows.

Consider the integral

$$I_C = \int_C e^{-t} (-t)^{M+a} (z-t)^{-M} dt$$

where $a \in \mathbb{C}$, M is a real parameter, and C is one of two loops C', C'' shown in the picture.

Picture 3. The contours C' , C'' .

We assume that at the point of C where t is negative we have $\arg(-t) = 0$ and $\arg(z-t) \in (0, \pi)$. We are interested in asymptotics of the integral as M tends to $+\infty$.

Lemma 10. *Assume that $\text{Im } z > 0$ and $\arg(-z) \in (-\pi, 0)$. Assume that $\arg M = 0$ and M tends to $+\infty$. Then*

$$I_{C'} = i\pi^{1/2}(-zM)^{(2a+1)/4} \exp(2(-zM)^{1/2} - z/2) (1 + O(M^{-1/2})),$$

$$I_{C''} = \pi^{1/2}(e^{2\pi i(M+a)} - 1)(-zM)^{(2a+1)/4} \exp(-2(-zM)^{1/2} - z/2 - \pi ia) (1 + O(M^{-1/2})).$$

To prove the lemma one changes the integration variable, $u = tM^{-1/2}$, then

$$I_C = M^{(a+1)/2} \int_C \exp(M^{1/2}(-u + z/u) + z^2/2u^2) (-u)^a du (1 + O(M^{-1/2})),$$

and the integral becomes a standard integral of the steepest descent method, whose asymptotics come from critical points of the function $-u + z/u$.

REFERENCES

- [FMTV] Felder, G., Markov, Y., Tarasov, V., Varchenko, A.: Differential equations compatible with KZ equations, *Math. Phys. Anal. Geom.* **3** (2000), no. 2, 139–177.
- [MTV] Markov, Y., Tarasov, V., Varchenko, A.: The determinant of a hypergeometric period matrix, *Houston J. of Math.* **24** (1998) no. 2, 197–220.
- [TV1] Tarasov, V., Varchenko, A.: Duality for Knizhnik-Zamolodchikov and dynamical equations, *Acta Appl. Math.* **73** (2002), no. 1-2, 141–154.
- [TV2] Tarasov, V., Varchenko, A.: Selberg type integrals associated with sl_3 , Preprint (2003) 1–11, [math.QA/0302148](#).